

## SUMMARY

The outlook May and for May–July is for river flows to be below normal for southern and central England and in the normal to below normal range elsewhere. Groundwater levels in May, and for the next three months, are likely to be in the normal to below normal range across the UK.

### Rainfall:

Rainfall in April was substantially below average for most of the UK with the exception of north-west Scotland. Large areas of southern and eastern England recorded less than a third of the average expected for the month.

The rainfall outlook (issued by the Met Office on 01.05.2022) for May and for the May–July period suggests near average rainfall is more likely to prevail than wetter or drier than average conditions.

### River flows

River flows in April were mainly below normal across the UK, notably or exceptionally so in south-west England, Wales, and western Scotland. There was an exception in the far north-west of Scotland where flows were above normal following above average rainfall in that area.


River flows in May are likely to show similar patterns to April. Below normal flows are expected across much of central, southern and eastern England, with normal to below normal flows being most likely across the rest of the country. A similar picture is seen for the May–July timeframe, albeit with a higher likelihood of normal flows in the north and west of the UK.

### Groundwater:

Groundwater levels in April were normal to below normal across England and Wales. Levels in south Wales and in some boreholes in southern England were below normal, exceptionally so in one case in south Wales.

Levels in May are likely to continue in the normal to below normal range at most boreholes in England and Wales. Below normal levels are likely in southern England. Levels for one borehole in the Scottish Borders are likely to be exceptionally high. The three-month outlook is very similar to the one-month outlook.

The Hydrological Outlook UK provides an outlook for the water situation for the UK over the next three months and beyond. For guidance on how to interpret the outlook, a wider range of information, and a full description of underpinning methods, please visit the website: [www.hydoutuk.net](http://www.hydoutuk.net)



River flows are most likely to be within the normal to below normal range for May–July in most of the UK

Groundwater levels are likely to be normal to below normal for May–July in most of the UK

River flows are likely to be below normal for May and over May–July for southern and central England

Shaded areas show principal aquifers

## About the Hydrological Outlook:

This document presents an outlook for the UK water situation for the next 1 – 3 months and beyond, using observational datasets, meteorological forecasts and a suite of hydrological modelling tools. The outlook is produced in a collaboration between the UK Centre for Ecology and Hydrology (UKCEH), British Geological Survey (BGS), the Met Office, the Environment Agency (EA), Natural Resources Wales (NRW), the Scottish Environment Protection Agency (SEPA), and for Northern Ireland, the Department for Infrastructure – Rivers (DfIR).

## Data and Models:

The Hydrological Outlook depends on the active cooperation of many data suppliers. This cooperation is gratefully acknowledged. Historic river flow and groundwater data are sourced from the UK National River Flow Archive and the National Groundwater Level Archive. Contemporary data are provided by the EA, SEPA, NRW and DfIR. These data are used to initialise hydrological models, and to provide outlook information based on statistical analysis of historical analogues.

Climate forecasts are produced by the Met Office. Hydrological modelling is undertaken by UKCEH using the Grid-to-Grid, PDM and CLASSIC hydrological models and by the EA using CATCHMOD. Hydrogeological modelling uses the R-groundwater model run by BGS and CATCHMOD run by the EA. Supporting documentation is available from the Outlooks website:

<https://www.hydoutuk.net/about/methods>

## Presentation:

The language used in the summary presented overleaf generally places flows and groundwater levels into just three classes, i.e. below normal, normal, and above normal. However, the underpinning methods use as many as seven classes as defined in the graphic to the right, i.e. the summary uses a simpler classification than some of the methods. On those occasions when it is appropriate to provide greater discrimination at the extremes the terminology and definitions of the seven class scheme will be adopted.

Percentile range of historic values for relevant month

Exceptionally high flow	> 95
Notably high flow	87-95
Above normal	72-87
Normal range	28-72
Below normal	13-28
Notably low flow	5-13
Exceptionally low flow	< 5

## Disclaimer and liability:

The Hydrological Outlook partnership aims to ensure that all Content provided is accurate and consistent with its current scientific understanding. However, the science which underlies hydrological and hydrogeological forecasts and climate projections is constantly evolving. Therefore any element of the Content which involves a forecast or a prediction should not be relied upon as though it were a statement of fact. To the fullest extent permitted by applicable law, the Hydrological Outlook Partnership excludes all warranties or representations (express or implied) in respect of the Content.

Your use of the Content is entirely at your own risk. We make no warranty, representation or guarantee that the Content is error free or fit for your intended use.

From April 2018 the Hydrological Outlook is supported by the Natural Environment Research Council funded [UK-SCAPE](#) and [Hydro-JULES](#) Programmes.

## Copyright:

Some of the features displayed on the maps contained in this report are based on the following data with permission of the controller of HMSO.

- (i) Ordnance Survey data. © Crown copyright and/or database right 2005. Licence no. 100017897.
- (ii) Land and Property Services data. © Crown copyright and database right, S&LA 145.
- (iii) Met Office rainfall data. © Crown copyright.

All rights reserved. Unauthorised reproduction infringes crown copyright and may lead to prosecution or civil proceedings.

## Further information:

For more detailed information about the Hydrological Outlook, and the derivation of the maps, plots and interpretation provided in this outlook, please visit the Hydrological Outlook UK website.

The website features a host of other background information, including a wider range of sources of information which are used in the preparation of this Outlook.

## Contact:

Hydrological Outlooks UK, UK Centre for Ecology & Hydrology, Wallingford, Oxfordshire, OX10 8BB  
t: 01491 692371 e: [enquiries@hydoutuk.net](mailto:enquiries@hydoutuk.net)

## Reference for the Hydrological Outlook:

Hydrological Outlook UK, 2022, May, UK Centre for Ecology and Hydrology, Oxfordshire UK, Online, <https://www.hydoutuk.net/latest-outlook/>

## Other Sources of Information:

The Hydrological Outlook should be used alongside other sources of up-to-date information on the current water resources status and flood risk.

Environment Agency Water Situation Reports: provides summary of water resources status on a monthly and weekly basis for England:

<https://www.gov.uk/government/collections/water-situation-reports-for-england>

Flood warnings are continually updated, and should be consulted for an up-to-date and localised assessment of flood risk:

Environment Agency: <https://flood-warning-information.service.gov.uk/map>

Natural Resources Wales: <https://flood-warning.naturalresources.wales/>

Scottish Environment Protection Agency: <https://www.sepa.org.uk/flooding.aspx>

Hydrological Summary for the UK: provides summary of current water resources status for the UK:

<https://nrfa.ceh.ac.uk/monthly-hydrological-summary-uk>

UK Met Office forecasts for the UK: <https://www.metoffice.gov.uk/#?tab=regionalForecast>

UK Water Resources Portal: monitor the UK hydrological situation in near real-time including rainfall, river flow, groundwater and soil moisture from COSMOS-UK:

<https://eip.ceh.ac.uk/hydrology/water-resources/>

# Monthly mean river flows simulated by the Grid-to-Grid hydrological model



Period: April 2022

Issue date: 05.05.2022

This map shows the simulated monthly mean flow across Great Britain for last month, ranked in terms of 54 years of historical flow estimates (1963 – 2016).

These flows are produced by the 1km resolution Grid-to-Grid (G2G) hydrological model, which is run up to the end of each calendar month using observed rainfall and MORECS potential evaporation as input.

Note that the G2G model provides estimates of natural flows.





# Current Daily Simulated Subsurface Water Storage Conditions

Based on subsurface water storage estimated for 30<sup>th</sup> April 2022

Issue date: 05.05.2022


These maps are based on Grid-to-Grid (G2G) hydrological model simulated subsurface water storage, expressed as an anomaly from the historical monthly mean. To highlight areas that are particularly wet or dry, the storage anomaly is presented here using a colour scale highlighting water storage relative to historical extremes. The maps below show the “relative wetness” which combines maps previously shown separately as the “relative wetness” and “relative dryness”.

These maps do not provide a forecast and are not maps of soil moisture. Instead they indicate areas which are particularly wet or dry. Rainfall in areas with high positive relative wetness could result in flooding in the coming days/weeks. Areas of negative relative wetness provide an indication of locations which are particularly dry, and little or no rain in these areas could potentially lead to (or prolong) a drought.





**SUMMARY:** At the end of April, subsurface water levels were generally lower (drier) than normal across Britain, very dry in many areas of central and southern England.

## Relative wetness

Water storage anomaly as a % of maximum (positive wetness) or minimum (negative wetness) storage anomaly (zero indicates average value)



Labels refer to estimated storage on *final day* of named month



\*Example month displaying extreme negative wetness

# Return Period of Rainfall Required to Overcome Dry Conditions

Period: May 2022 – Oct 2022

Issue date: 05.05.2022

These maps show the **return period** of the rainfall required to overcome dry conditions simulated using the Grid-to-Grid (G2G) hydrological model. The maps are coloured according to the return period of accumulated rainfall required to overcome the estimated current subsurface water storage deficit over the next few months.

These maps do not provide a drought forecast. Instead they indicate the return period of rainfall required to overcome the dry conditions for the following 6 months based on current conditions.

**SUMMARY:** During **May to July**, regions in southern and eastern England would require rainfall with a return period of between 5 and 50 years to overcome the dry conditions. Elsewhere, not particularly unusual rainfall (<5 year return periods) would be required to return to average conditions for this time of year.

During **August to October**, Great Britain will not require particularly unusual rainfall (<5 year return periods) to return to average conditions for the time of year.



## SCOTLAND


HR Highlands Region  
NER North East Region  
TR Tay Region  
FR Forth Region  
CR Clyde Region  
TWR Tweed Region  
SR Solway Region

## ENGLAND

N Northumbria  
NW North West  
Y Yorkshire  
ST Severn Trent  
A Anglian  
T Thames  
S Southern  
W Wessex  
SW South West








## WALES

WEL Welsh



## NORTHERN IRELAND

This method cannot currently be used in Northern Ireland

Rainfall amount / Probability		Return period (years)	
Low (this rain is likely to occur)	> 20%		< 5
	< 20%		5 - 10
	< 10%		10 - 25
	< 4%		25 - 50
High (less likely)	< 2%		50 - 100
	< 1%		100 - 200
Extreme (unlikely but still possible)	< 0.5%		> 200

# Estimate of Additional Rainfall Required to Overcome Dry Conditions

Based on subsurface water storage estimated for 30<sup>th</sup> April 2022

Issue date: 05.05.2022

These maps show the Grid-to-Grid (G2G) hydrological model simulated subsurface water storage, expressed as an anomaly from the historical monthly mean (1981-2010), presented on a 1km grid and as regional means.

**Subsurface storage deficits**, i.e. where the subsurface water storage anomaly is less than zero, are highlighted by the red/pink colours.

The **subsurface storage deficit (mm)** can be interpreted as an estimate of additional rainfall that would be required in future months to overcome dry conditions (i.e. rainfall in addition to what is expected on average). Regional mean values of additional rainfall required are provided in the table below.

*Regional estimate of additional rainfall required (mm)*

## SCOTLAND




21	HR	Highlands Region
11	NER	North East Region
23	TR	Tay Region
27	FR	Forth Region
20	CR	Clyde Region
22	TWR	Tweed Region
25	SR	Solway Region

## ENGLAND


17	N	Northumbria
26	NW	North West
25	Y	Yorkshire
39	ST	Severn Trent
49	A	Anglian
54	T	Thames
61	W	Wessex
63	S	Southern
49	SW	South West

## WALES

36	WEL	Welsh
----	-----	-------




*Water storage deficit  
(anomaly, mm)*





**SUMMARY:** The outlook for May and for May to July is for below normal flows in central and southern England, and normal to below normal flows in northern England and Northern Ireland. In north-east Scotland flows are likely to be normal to above normal. Please note there are not many forecasts available for north-west Britain.

River flow outlook for May 2022




1-month flow outlook

Outlooks from hydrological analogues are based on a comparison of river flow during recent months with flows during the same months in previous years at a set of approximately 90 sites from across the UK. These sites are depicted on the two maps. Years with observed flows that most closely resemble current conditions are identified as the best analogues and the outlook is based on extrapolating from current conditions based on these analogues.

It is, however, often the case that a simpler forecast based on the persistence of river flow provides a better forecast than provided by analogy. This is particularly true for slowly responding catchments associated with aquifer outcrops.

Both methods are considered at each site and the forecast from the method with the higher confidence is presented. A simple classification of flows is used (high, medium and low) as indicated by the colours of the dots, with the confidence

River flow outlook for May - Jul 2022



3-month flow outlook





of the forecast being represented by the size of the dot. A tag on the dot indicates which method has been used in each instance.

Period: May 2022




These figures provide insight into the hydrological analogue methodology for a set of sites from across the UK.

In each of the time series graphs the bold black line represents the observed flow during the past six months. The grey band indicates the normal flow range (the normal band includes 44%

of observed flows in each month). The selected analogues are shown as thin lines and the trajectories that flows took in the following month are also shown. The forecast is shown as the dashed red line, and in each plot it states whether this has come from the analogues or has been generated on the basis of persistence.



Issued on 05.05.2022 using data to the end of April 2022



No forecast available







Period: May 2022 – July 2022




These figures provide insight into the hydrological analogue methodology for a set of sites from across the UK.

In each of the time series graphs the bold black line represents the observed flow during the past nine months. The grey band indicates the normal flow range (the normal band includes 44%

of observed flows in each month). The selected analogues are shown as thin lines and the trajectories that flows took in the following three months are also shown. The forecast is shown as the dashed red line, and in each plot it states whether this has come from the analogues or has been generated on the basis of persistence.



Issued on 05.05.2022 using data to the end of May 2022




No forecast available

Period: May 2022 – October 2022


Issued on 03.05.2022 using data to the end of April 2022

The outlook for May indicates that flows are most likely to be below normal for southern and central England, normal for Northern Ireland, and normal to below normal for the rest of the UK. The April-May-June outlook indicates that conditions are likely to persist for southern and central England, and Wales, with the rest of the UK likely to experience normal flows.


### 1-month river flow outlook starting May 2022



### 3-month river flow outlook starting May 2022



### 6-month river flow outlook starting May 2022



This outlook is based on monthly ensembles of historical sequences of observed climate (rainfall and potential evapotranspiration) that form input to a hydrological model. The outputs are probabilistic simulations of the average river flow over the forecast period (1 to 12 months ahead), at each location. The simulations are generated by the GR4J conceptual rainfall-runoff model from IRSTEA (France) calibrated on observed or naturalised flows.

The bar plot maps show the outlook distribution for 1, 3 and 6-month period for 64 catchments across England and Wales. Each bar plot represents the probabilistic distribution of the simulated river flow compared to the historical river flow, for the same n-month period. The probabilities fall within five categories, classified as: low, below normal, normal, above normal and high.

This outlook is based entirely on historical sequences and therefore does not contain any knowledge of the state of the atmosphere and ocean. It is hence possible that some of the historical sequences used might be inconsistent with current large-scale atmospheric conditions and would therefore be unlikely to occur in the next few months.



This outlook is based on monthly ensembles of historical sequences of observed climate (rainfall and potential evapotranspiration) that form input to a hydrological model. The outputs are probabilistic simulations of the average river flow over the forecast period (1 to 12 months ahead), at each location. The simulations are generated by the GR4J conceptual rainfall-runoff model from IRSTEA (France) calibrated on observed or naturalised flows.

The stack diagrams show the variation over time of the outlook distribution for a number of individual catchments. Each graph represents variation over time of the number of simulated river flows, in each month ensemble, that fall within each of seven categories: exceptionally low, notably low, below normal, normal, above normal, notably high and exceptionally high. The categories represent cumulative flow conditions, e.g. For 3-month, the simulated total 3-month flow compared to the historical 3-month flow distribution. The monthly variations can be compared to the long-term average distribution of river flows (shown as columns

on the right of each timeline graph).

This outlook is based entirely on historical sequences and therefore does not contain any knowledge of the state of the atmosphere and ocean. It is hence possible that some of the historical sequences used might be inconsistent with current large-scale atmospheric conditions and would therefore be unlikely to occur in the next few months.



**SUMMARY:** During May, river flows in most regions are likely to be in the *Normal range* or below. Anglian Region is most likely to be *Below normal*.

**Over the next 3 months** river flows in southern England are likely to be *Below normal* or lower. Elsewhere, river flows are likely to be in the *Normal range* or below.

These forecasts are produced by using five members of the Met Office rainfall forecast ensemble as input to a water balance hydrological model to provide the five estimates of river flows shown on the left for one month and three months ahead.

Regional forecast monthly-mean river flows are derived from the average of 1km river flow estimates within each region and ranked in terms of 54 years of historical flow estimates (1963 – 2016).

The five maps illustrate the wide range of possible flows and while there is a 50% chance of flows between the 1<sup>st</sup> and 3<sup>rd</sup> quartiles, actual flows may be more extreme than the flows derived using the highest or lowest rainfall forecasts.






Lowest rainfall forecast

1<sup>st</sup> quartile

Median

3<sup>rd</sup> quartile

Highest rainfall forecast



## Key

Exceptionally high flow  
Notably high flow  
Above normal  
Normal range  
Below normal  
Notably low flow  
Exceptionally low flow

Percentile range of historic values for relevant month

> 95
87-95
72-87
28-72
13-28
5-13
< 5






Lowest rainfall forecast

1<sup>st</sup> quartile

Median

3<sup>rd</sup> quartile

Highest rainfall forecast



## SCOTLAND


HR Highlands Region  
NER North East Region  
TR Tay Region  
FR Forth Region  
CR Clyde Region  
TWR Tweed Region  
SR Solway Region

## ENGLAND

N Northumbria  
NW North West  
Y Yorkshire  
ST Severn Trent  
A Anglian  
T Thames  
S Southern  
W Wessex  
SW South West

## WALES

WEL Welsh





**NORTHERN IRELAND**  
This method cannot currently be used in Northern Ireland

The regional maps illustrating the regional river flows for five members of the Met Office ensemble of rainfall forecasts give some indication of the range of possible river flows in the coming months. As noted previously, the actual flows could be more extreme than the flows generated by either the lowest or highest members of the rainfall ensemble.

The bar charts (below) give further insight into the range of river flow forecasts by considering all members of the forecast rainfall ensemble. The regional bar charts show the percentage of ensemble forecasts falling in each of the flow categories as generated by the monthly-resolution water-balance model. As before results are averaged by region then ranked in terms of 54 years of historical regional flow estimates (1963 – 2016).

**SUMMARY:** During May, river flows in most regions are likely to be in the *Normal range* or below. Anglian Region is most likely to be *Below normal*.

**Over the next 3 months** river flows in southern England are likely to be *Below normal* or lower. Elsewhere, river flows are likely to be in the *Normal range* or below.



## SCOTLAND

HR Highlands Region  
NER North East Region  
TR Tay Region  
FR Forth Region  
CR Clyde Region  
TWR Tweed Region  
SR Solway Region

## ENGLAND

N Northumbria  
NW North West  
Y Yorkshire  
ST Severn Trent  
A Anglian  
T Thames  
S Southern  
W Wessex  
SW South West

## WALES

WEL Welsh



## NORTHERN IRELAND

This method cannot currently be used in Northern Ireland

The maps illustrating the regional river flows for five members of the Met Office ensemble of rainfall forecasts give some indication of the range of possible river flows in the coming months. As noted previously, the actual flows could be more extreme than the flows generated by either the lowest or highest members of the rainfall ensemble.

The tables below give further insight into the range of river flow forecasts by considering all members of the forecast rainfall ensemble. The numbers in the tables are the percentage of ensemble forecasts falling in each of the flow categories as generated by the monthly-resolution water-balance model. As before results are averaged by region then ranked in terms of 54 years of historical regional flow estimates (1963 – 2016).

**SUMMARY:** During May, river flows in most regions are likely to be in the *Normal range* or below. Anglian Region is most likely to be *Below normal*.

**Over the next 3 months** river flows in southern England are likely to be *Below normal* or lower. Elsewhere, river flows are likely to be in the *Normal range* or below.

## SCOTLAND

HR Highlands Region  
NER North East Region  
TR Tay Region  
FR Forth Region  
CR Clyde Region  
TWR Tweed Region  
SR Solway Region

## ENGLAND

N Northumbria  
NW North West  
Y Yorkshire  
ST Severn Trent  
A Anglian  
T Thames  
S Southern  
W Wessex  
SW South West

## WALES

WEL Welsh



## NORTHERN IRELAND

This method cannot currently be used in Northern Ireland

1-month ahead	A	NW	N	ST	SW	S	T	Welsh	W	Y	CR	FR	HR	NER	SR	TR	TWR
Exceptionally high flow	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Notably high flow	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Above normal	2	5	5	2	0	0	0	5	0	2	0	2	7	7	5	2	2
Normal range	36	57	52	48	57	50	50	50	43	50	57	52	48	45	62	55	55
Below normal	55	12	26	24	19	26	26	24	26	26	21	21	14	21	14	21	21
Notably low flow	7	26	17	10	14	24	24	14	31	21	21	17	26	24	19	21	21
Exceptionally low flow	0	0	0	17	10	0	0	7	0	0	0	7	5	2	0	0	0

3-months ahead	A	NW	N	ST	SW	S	T	Welsh	W	Y	CR	FR	HR	NER	SR	TR	TWR
Exceptionally high flow	0	0	0	0	0	0	0	0	0	0	0	0	5	0	0	0	0
Notably high flow	0	0	5	0	0	0	0	5	0	5	5	5	0	5	0	0	5
Above normal	0	5	0	5	5	0	0	0	0	0	0	0	2	0	7	5	0
Normal range	14	45	60	26	33	14	12	38	14	52	43	60	43	55	64	40	64
Below normal	83	24	21	55	43	45	81	29	55	24	33	21	24	24	14	36	17
Notably low flow	2	10	12	7	12	38	7	21	24	12	12	7	10	10	7	12	7
Exceptionally low flow	0	17	2	7	7	2	0	7	7	7	7	7	17	7	7	7	7




Period: May 2022 – July 2022

Issued on 10.05.2022 using data to the end of April


Over the next month, normal to notably low groundwater levels are forecast across most of England and Wales. There are a few exceptions, including exceptionally low groundwater levels forecast at the Beeding Hill site in the southern Chalk in the one month forecast, tending towards below normal over three months. Note there are a reduced number of modelled sites. This is due to Covid-19 restrictions on access to sites in England and IT issues in Scotland.

These forecasts are produced by running five members of the Met Office ensemble climate forecast through groundwater models of observation borehole hydrographs at 42 sites across the country. The sites are distributed across the principal aquifers.

Based on the distribution of observed historical groundwater levels in a given month, seven categories have been derived for each site: very low, low, below normal, normal, above normal, high, and very high. The forecast groundwater level is assigned to one of these seven categories depending on where it falls within the distribution of the historically observed values.




1-month outlook



3-month outlook

Key	Percentile range of historic observed values for relevant month
Exceptionally high levels	> 95
Notably high levels	87-95
Above normal	72-87
Normal	28-72
Below normal	12-28
Notably low levels	5-13
Exceptionally low levels	< 5






The Hydrological Outlook UK provides an outlook for the water situation for the UK over the next three months and beyond. For guidance on how to interpret the outlook, a wider range of information, and a full description of underpinning methods, please visit the website: [www.hydoutuk.net](http://www.hydoutuk.net)

# Outlook based on modelled groundwater from historical climate

Period: May 2022 – April 2023

Issued on 10.05.2022 using data to the end of April

At Little Bucket Farm, groundwater levels are predicted to remain below normal over the next 6 months, with normal to below normal levels predicted from 6 to 12 months. Elsewhere in the Chalk groundwater levels are predicted to be predominantly normal over the next 12 months. In the Permo-Triassic sandstone at Bussels, normal to below normal levels are predicted to prevail over the next 12 months, while at Skirwith below normal to notably low levels are predicted over the next 6 months, tending towards normal to above normal levels from 6 to 12 months.



This outlook is based on monthly ensembles of historical sequences of observed climate (rainfall and potential evapotranspiration) that form input to hydrological models. The outputs are probabilistic simulations of the average groundwater level over the forecast horizon (3 to 12 months ahead), at each location.

The graphs show variation over time of the number of simulated groundwater levels in each monthly ensemble,

that fall within each the seven categories: exceptionally low, notably low, below normal, normal, above normal, notably high and exceptionally high. The monthly variations can be compared to the long-term average distribution of levels, which are shown as columns on the left and right of each graph.

This outlook is based entirely on historical sequences and therefore does not contain any knowledge of the state of

the atmosphere and ocean. It is hence possible that some of the historical sequences used might be inconsistent with current large-scale atmospheric conditions and would therefore be unlikely to occur in the next few months.