


Monthly mean river flows simulated by the Grid-to-Grid hydrological model

Period: March 2023

This map shows the simulated monthly mean flow across Great Britain for last month, ranked in terms of 54 years of historical flow estimates (1963 - 2016).

These flows are produced by the 1km resolution Grid-to-Grid (G2G) hydrological model, which is run up to the end of each calendar month using observed rainfall and MORECS potential evaporation as input.

Note that the G2G model provides estimates of natural flows.



pixel ranked in terms of historic % flow estimates (1963-2016)

Exceptionally high flow Notably high flow 87-95 Above normal 72-87 Normal range 28-72 Below normal 13-28 Notably low flow 5-13 Exceptionally low flow

The Hydrological Outlook UK provides an outlook for the water situation for the UK over the next three months and beyond. For guidance on how to interpret the outlook, a wider range of information, and a full description of underpinning methods, please visit the website: www.hydoutuk.net

Issue date: 02.03.2023




Current Daily Simulated Subsurface Water Storage Conditions

Based on subsurface water storage estimated for 28 February 2023

These maps are based on Grid-to-Grid (G2G) hydrological model simulated subsurface water storage, expressed as an anomaly from the historical monthly mean. To highlight areas that are particularly wet or dry, the storage anomaly is presented here using a colour scale highlighting water storage relative to historical extremes. The maps below show the "relative wetness" which combines maps previously shown separately as the "relative wetness" and "relative dryness".

These maps do not provide a forecast and are not maps of soil moisture. Instead they indicate areas which are particularly wet or dry. Rainfall in areas with high positive relative wetness could result in flooding in the coming days/weeks. Areas of negative relative wetness provide an indication of locations which are particularly dry, and little or no rain in these areas could potentially lead to (or prolong) a drought.

SUMMARY: At the end of February, subsurface water levels were lower (drier) than normal in most of Great Britain.



*Example month displaying extreme negative wetness

The Hydrological Outlook UK provides an outlook for the water situation for the UK over the next three months and beyond. For guidance on how to interpret the outlook, a wider range of information, and a full description of underpinning methods, please visit the website: www.hydoutuk.net




Return Period of Rainfall Required to Overcome Dry Conditions

Period: March 2023 - August 2023

These maps show the return period of the rainfall required to overcome dry conditions simulated using the Grid-to-Grid (G2G) hydrological model. The maps are coloured according to the return period of accumulated rainfall required to overcome the estimated current subsurface water storage deficit over the next few months.

These maps do not provide a drought forecast. Instead they indicate the return period of rainfall required to overcome the dry conditions for the following 6 months based on current conditions.

SUMMARY: During March most of Great Britain will require rainfall with a return period of between 5 – 25 years to return to normal. The Anglian region requires an additional 3 months of rainfall with a return period of 5 - 10 years to return to average conditions for the time of year.




SCOTLAND

HR Highlands Region
NER North East Region
TR Tay Region
FR Forth Region
CR Clyde Region
TWR Tweed Region
SR Solway Region
SR Solway Region

NORTHERN IRELAND

Issue date: 02.03.2023

This method cannot currently be used in Northern Ireland



The Hydrological Outlook UK provides an outlook for the water situation for the UK over the next three months and beyond. For guidance on how to interpret the outlook, a wider range of information, and a full description of underpinning methods, please visit the website: www.hydoutuk.net



Estimate of Additional Rainfall Required to Overcome Dry Conditions

Based on subsurface water storage estimated for 28 February 2023

These maps show the Grid-to-Grid (G2G) hydrological model simulated subsurface water storage, expressed as an anomaly from the historical monthly mean (1981-2010), presented on a 1km grid and as regional means.

Subsurface storage deficits, i.e. where the subsurface water storage anomaly is less than zero, are highlighted by the red/pink colours.

The subsurface storage deficit (mm) can be interpreted as an estimate of additional rainfall that would be required in future months to overcome dry conditions (i.e. rainfall in addition to what is expected on average). Regional mean values of additional rainfall required are provided in the table below.

Regional estimate of additional rainfall required (mm)

SCOTLAND


- 11 HR Highlands Region
- 9 NER North East Region
- 12 TR Tay Region
- 12 FR Forth Region
- 12 CR Clyde Region
- 11 TWR Tweed Region
- 14 SR Solway Region

ENGLAND

- 5 N Northumbria
- 14 NW North West
- 19 Y Yorkshire
- 24 ST Severn Trent
- 39 A Anglian
- 32 T Thames
- 17 W Wessex
- 5 S Southern
- 26 SW South West


WALES

26 WEL Welsh





Issue date: 02.03.2023



The Hydrological Outlook UK provides an outlook for the water situation for the UK over the next three months and beyond. For guidance on how to interpret the outlook, a wider range of information, and a full description of underpinning methods, please visit the website: www.hydoutuk.net