

SUMMARY The river flow outlook for January favours normal flows across the UK, although above normal flows may persist in parts of central and southern England that have seen the highest rainfall in recent weeks. In contrast, January – March outlook favours wetter conditions that are more likely to manifest themselves in normal to above normal flows in the north and west, with normal flows most likely elsewhere. While groundwater is a more mixed picture (reflecting both recent rainfall patterns and aquifer properties), normal to above normal levels are expected across the UK over the January – March period.

Rainfall:


The far north of Scotland saw notably above average December rainfall, whereas southern Scotland and Northern Ireland were relatively dry. Similarly, parts of northern England and north Wales saw above average rainfall, whereas further south it was drier than average, particularly in the far south. The forecast (issued by Met Office on 23.12.2024) indicates the chance of a dry January is slightly higher than normal. In contrast, January-March has a higher-than-normal chance of being wet due to an increased likelihood of westerly winds.

River flows:

River flows in December were notably or exceptionally high in the far north of Scotland, with new records for December in some catchments. Elsewhere, river flows were predominantly in the normal range or above normal, with above normal flows most prevalent across central England and north Wales. The outlook for January is for river flows to be in the normal range across most of the UK, but above normal flows may persist in parts of central England and the south, where early January has already seen significant rainfall. The January-March outlook favours flows being normal to above normal in northern Britain and in the normal range further south.

Groundwater:

Groundwater levels in December were in the normal range in the far south of England, and normal to above normal in most boreholes in central and northern England. Below normal levels were registered at some sites in central Scotland and Northern Ireland. The January Outlook is for a similar picture of normal to above normal levels predominating across the UK, with above normal levels most likely in central areas (e.g. the Chilterns and Jurassic Limestone), and normal levels further south. Over the January-March period, some areas will see levels receding into the normal range, while others will see increases due to the delayed impact of recent rainfall – hence, the outlook is for normal to above normal levels at the national scale.



Groundwater levels for the UK are likely to be normal to above normal over the January to March period.

River flows in northern Britain are likely to be in the normal range in January, and normal to above normal in the January-March period.

River flows in southern Britain are likely to be in the normal range in January-March, although above normal flows may persist through January in some areas.

Shaded areas show principal aquifers

The UK Hydrological Outlook provides an outlook for the water situation for the United Kingdom over the next three months and beyond. For guidance on how to interpret the outlook, a wider range of information, and a full description of underpinning methods, please visit the website: www.hyoutuk.net

About the UK Hydrological Outlook:

This document presents an outlook for the UK water situation for the next 1-3 months and beyond, using observational datasets, meteorological forecasts and a suite of hydrological modelling tools. The outlook is produced in a collaboration between the UK Centre for Ecology & Hydrology (UKCEH), British Geological Survey (BGS), the Met Office, the Environment Agency (EA), Natural Resources Wales (NRW), the Scottish Environment Protection Agency (SEPA), and for Northern Ireland, the Department for Infrastructure – Rivers (DfIR).

Data and Models:

The UK Hydrological Outlook depends on the active cooperation of many data suppliers. This cooperation is gratefully acknowledged. Historic river flow and groundwater data are sourced from the [UK National River Flow Archive](#) and the [National Groundwater Level Archive](#). Contemporary data are provided by the EA, SEPA, NRW and DfIR. These data are used to initialise hydrological models, and to provide outlook information based on statistical analysis of historical analogues.

Climate forecasts are produced by the Met Office. Hydrological modelling is undertaken by UKCEH using the Grid-to-Grid and GR6J hydrological models. Hydrogeological modelling uses the AquilMod model run by BGS. Supporting documentation is available from the Outlooks website: <https://hydoutuk.net/about/methods>

Presentation:

The language used in the summary presented overleaf generally places flows and groundwater levels into just three classes, i.e. below normal, normal, and above normal. However, the underpinning methods use as many as seven classes as defined in the graphic to the right, i.e. the summary uses a simpler classification than some of the methods. On those occasions when it is appropriate to provide greater discrimination at the extremes the terminology and definitions of the seven class scheme will be adopted.

	Percentile range of historic values for relevant month
Exceptionally high flow	> 95
Notably high flow	87-95
Above normal	72-87
Normal range	28-72
Below normal	13-28
Notably low flow	5-13
Exceptionally low flow	< 5

Disclaimer and liability:

The UK Hydrological Outlook partnership aims to ensure that all Content provided is accurate and consistent with its current scientific understanding. However, the science which underlies hydrological and hydrogeological forecasts and climate projections is constantly evolving. Therefore any element of the Content which involves a forecast or a prediction should not be relied upon as though it were a statement of fact. To the fullest extent permitted by applicable law, the UK Hydrological Outlook Partnership excludes all warranties or representations (express or implied) in respect of the Content.

Your use of the Content is entirely at your own risk. We make no warranty, representation or guarantee that the Content is error free or fit for your intended use.

The UK Hydrological Outlook is supported by the Natural Environment Research Council funded NC-UK (NE/Y006208/1) and [Hydro-JULES](#) (NE/S017380/1) Programmes.

Copyright:

Some of the features displayed on the maps contained in this report are based on the following data with permission of the controller of HMSO.

- Ordnance Survey data. © Crown copyright and/or database right 2005. Licence no. 100017897.
- Land and Property Services data. © Crown copyright and database right, S&LA 145.
- Met Office rainfall data. © Crown copyright.
- The three month outlook contained in the hydrological outlook is licensed under the terms of the [Open Government Licence](#)

All rights reserved. Unauthorised reproduction infringes crown copyright and may lead to prosecution or civil proceedings.

Further information:

For more detailed information about the UK Hydrological Outlook, and the derivation of the maps, plots and interpretation provided in this outlook, please visit the UK Hydrological Outlook website. The website features a host of other background information, including a wider range of sources of information which are used in the preparation of this Outlook. Dynamic access to many of the outputs of the UK Hydrological Portal are available on the [UK Hydrological Outlooks Portal](#).

Contact:

UK Hydrological Outlooks, UK Centre for Ecology & Hydrology, Wallingford, Oxfordshire, OX10 8BB
t: 01491 838800 e: <https://hydoutuk.net/contact>

Reference for the UK Hydrological Outlook:

UK Hydrological Outlook, 10 January 2025, UK Centre for Ecology & Hydrology, Oxfordshire UK, Online, <https://www.hydoutuk.net/latest-outlook/>

Other Sources of Information:

The UK Hydrological Outlook should be used alongside other sources of up-to-date information on the current water resources status and flood risk.

Environment Agency Water Situation Reports: provides summary of water resources status on a monthly and weekly basis for England: <https://www.gov.uk/government/collections/water-situation-reports-for-england>

Flood warnings are continually updated, and should be consulted for an up-to-date and localised assessment of flood risk:

- Environment Agency: <https://flood-warning-information.service.gov.uk/map>
- Natural Resources Wales: <https://flood-warning.naturalresources.wales/>
- Scottish Environment Protection Agency: <https://www.sepa.org.uk/flooding.aspx>

Hydrological Summary for the UK: provides summary of current water resources status for the UK: <https://nfa.ceh.ac.uk/monthly-hydrological-summary-uk>

UK Met Office forecasts for the UK: <https://www.metoffice.gov.uk/>

UK Water Resources Portal: monitor the UK hydrological situation in near real-time including rainfall, river flow, groundwater and soil moisture from COSMOS-UK: <https://eip.ceh.ac.uk/hydrology/water-resources/>


December's mean river flows simulated by the Grid-to-Grid hydrological model

Issue date: 06.01.2025

This map shows the simulated monthly mean flow across Great Britain for last month, ranked in terms of 54 years of historical flow estimates (1963 – 2016).

These flows are produced by the 1km resolution Grid-to-Grid (G2G) hydrological model, which is run up to the end of each calendar month using observed rainfall and MORECS potential evaporation as input.

Note that the G2G model provides estimates of natural flows.



Current Daily Simulated Subsurface Water Storage Conditions

Based on subsurface water storage estimated for 31 December 2024



Issue date: 06.01.2025

These maps are based on Grid-to-Grid (G2G) hydrological model simulated subsurface water storage (water in the soil and groundwater), expressed as an anomaly from the historical monthly mean. To highlight areas that are particularly wet or dry, the storage anomaly is presented relative to historical extremes. Rainfall in WET areas with high positive relative wetness could result in flooding in the coming days/weeks. Areas of negative relative wetness indicate locations which are particularly DRY, and little or no rain in these areas could potentially lead to (or prolong) a drought. Maps of soil moisture only are available on the next page.

SUMMARY: Subsurface water stores are high (very wet) in most of Scotland, the Northwest of England and Wales. In the rest of the country, subsurface water stores are typical to wet for the time of year.

Relative wetness

Water storage anomaly as a % of maximum (positive wetness) or minimum (negative wetness) storage anomaly (zero indicates average value)



*Example month displaying extreme positive wetness

January 2025



CURRENT CONDITIONS

These maps are based on Grid-to-Grid (G2G) hydrological model simulated soil moisture, expressed as an anomaly from the historical monthly mean. To highlight areas that are particularly wet or dry, the soil moisture anomaly is presented relative to historical extremes. These maps are not a forecast; rather an indication of current conditions. Soil moisture will often look similar to total storage (shown on the previous slide), since total storage comprises both soil moisture and storage in the saturated zone.

SUMMARY: Soil moisture stores are high (very wet) in most of Scotland, the Northwest of England and Wales. In the rest of the country, soil moisture stores are typical to wet for the time of year.

Relative wetness

Soil moisture anomaly as a % of maximum (positive wetness) or minimum (negative wetness) storage anomaly (zero indicates average value)



*Example month displaying extreme positive wetness

Estimate of Additional Rainfall Required to Overcome Dry Conditions

Based on subsurface water storage estimated for 31 December 2024

Issue date: 06.01.2025

These maps show the Grid-to-Grid (G2G) hydrological model simulated subsurface water storage, expressed as an anomaly from the historical monthly mean (1981-2010), presented on a 1km grid and as regional means. Subsurface storage deficits, i.e. where the subsurface water storage anomaly is less than zero, are highlighted in red/pink.

The subsurface storage deficit (mm) can be interpreted as an estimate of additional rainfall that would be required in future months to overcome dry conditions (i.e. rainfall in addition to what is expected on average). Regional mean values of additional rainfall required are provided in the table below.

Regional estimate of additional rainfall required (mm)

SCOTLAND



- 0 HR Highlands Region
- 0 NER North East Region
- 0 TR Tay Region
- 0 FR Forth Region
- 0 CR Clyde Region
- 0 TWR Tweed Region
- 0 SR Solway Region

ENGLAND


- 0 N Northumbria
- 0 NW North West
- 0 Y Yorkshire
- 0 ST Severn Trent
- 0 A Anglian
- 0 T Thames
- 0 W Wessex
- 0 S Southern
- 9 SW South West

WALES

- 0 WEL Welsh



Water storage deficit (anomaly; mm)




Return Period of Rainfall Required to Overcome Dry Conditions

Period: January 2025 - June 2025




Issue date: 06.01.2025

These maps show the return period of the rainfall required to overcome dry conditions simulated using the Grid-to-Grid (G2G) hydrological model. The maps are coloured according to the return period of accumulated rainfall required to overcome the estimated current subsurface water storage deficit over the next one to six months (areas with no storage deficit will always be white). These maps do not provide a drought forecast; instead they indicate whether particularly heavy rainfall would be required to return to normal conditions for the time of year.

SUMMARY: The South West has a small subsurface storage deficit but does not need an unusual amount of rainfall to recover.




- SCOTLAND**
 HR Highlands Region
 NER North East Region
 TR Tay Region
 FR Forth Region
 CR Clyde Region
 TWR Tweed Region
 SR Solway Region
- ENGLAND**
 N Northumbria
 NW North West
 Y Yorkshire
 ST Severn Trent
 A Anglian
 T Thames
 S Southern
 W Wessex
 SW South West
- WALES**
 WEL Welsh



SUMMARY:

The January outlook indicates river flows across most of the UK are likely to be normal to above normal. In the south-west of England, flows in some catchments are likely to be normal to below normal. Over the January–March time frame, this pattern is likely to persist.


River flow outlook for Jan 2025



1-month flow outlook

Outlooks from hydrological analogues are based on a comparison of river flow during recent months with flows during the same months in previous years at a set of approximately 90 sites from across the UK. These sites are depicted on the two maps. Years with observed flows that most closely resemble current conditions are identified as the best analogues and the outlook is based on extrapolating from current conditions based on these analogues.

River flow outlook for Jan - Mar 2025



3-month flow outlook

It is, however, often the case that a simpler forecast based on the persistence of river flow provides a better forecast than provided by analogy. This is particularly true for slowly responding catchments associated with aquifer outcrops.

Both methods are considered at each site and the forecast from the method with the higher confidence is presented. A simple classification of flows is used (high, medium and low) as indicated by the colours of the dots, with the confidence

of the forecast being represented by the size of the dot. A tag on the dot indicates which method has been used in each instance.









Period: January 2025

Issued on 08.01.2025 using data to the end of December 2024

These figures provide insight into the hydrological analogue methodology for a set of sites from across the UK.

In each of the time series graphs the bold black line represents the observed flow during the past six months. The grey band indicates the normal flow range (the normal band includes 44%

of observed flows in each month). The selected analogues are shown as thin lines and the trajectories that flows took in the following month are also shown. The forecast is shown as the dashed red line, and in each plot it states whether this has come from the analogues or has been generated on the basis of persistence.






Period: January 2025 – March 2025

Issued on 08.01.2025 using data to the end of December 2024

These figures provide insight into the hydrological analogue methodology for a set of sites from across the UK.

In each of the time series graphs the bold black line represents the observed flow during the past nine months. The grey band indicates the normal flow range (the normal band includes 44%

of observed flows in each month). The selected analogues are shown as thin lines and the trajectories that flows took in the following three months are also shown. The forecast is shown as the dashed red line, and in each plot it states whether this has come from the analogues or has been generated on the basis of persistence.




Period: January 2025 – June 2025


Issued on 07.01.2025 using data to the end of December 2024

The outlook for January indicates that flows are likely to be normal to below normal across southwestern England and East Anglia. River flows in northern England and northeastern Scotland are likely to be normal to above normal. Elsewhere in the UK, flows are likely to be in the normal range. The January to March outlook indicates that flows across the UK are likely to be in the normal range.


1-month river flow outlook starting Jan 2025



3-month river flow outlook starting Jan 2025



6-month river flow outlook starting Jan 2025





This outlook is based on monthly ensembles of historical sequences of observed climate (rainfall and potential evapotranspiration) that form input to a hydrological model. The outputs are probabilistic simulations of the average river flow over the forecast period (1 to 12 months ahead), at each location. The simulations are generated by the GR6J conceptual rainfall-runoff model from INRAE (France) calibrated on observed or naturalised flows.

The bar plot maps show the outlook distribution for 1, 3 and 6-month period for 64 catchments across England and Wales. Each bar plot represents the probabilistic distribution of the simulated river flow compared to the historical river flow, for the same n-month period. The probabilities fall within five categories, classified as: low, below normal, normal, above normal and high.

This outlook is based entirely on historical sequences and therefore does not contain any knowledge of the state of the atmosphere and ocean. It is hence possible that some of the historical sequences used might be inconsistent with current large-scale atmospheric conditions and would therefore be unlikely to occur in the next few months.

Please note that *Outlooks based on modelled flow from historical climate* from October 2023 onwards were generated using GR6J model, whereas until September 2023, they were produced using GR4J model. For more details, please see the section on River flow from historical climate at this link: <https://hydoutuk.net/about/methods/river-flows>



This outlook is based on monthly ensembles of historical sequences of observed climate (rainfall and potential evapotranspiration) that form input to a hydrological model. The outputs are probabilistic simulations of the average river flow over the forecast period (1 to 12 months ahead), at each location. The simulations are generated by the GR6J conceptual rainfall-runoff model from INRAE (France) calibrated on observed or naturalised flows.

The stack diagrams show the variation over time of the outlook distribution for a number of individual catchments. Each graph represents variation over time of the number of simulated river flows, in each month ensemble, that fall within each of seven categories: exceptionally low, notably low, below normal, normal, above normal, notably high and exceptionally high. The categories represent cumulative flow conditions, e.g. For 3-month, the simulated total 3-month flow compared to the historical 3-month flow distribution. The monthly variations can be compared to the long-term average distribution of river flows (shown as columns on

the right of each timeline graph).

This outlook is based entirely on historical sequences and therefore does not contain any knowledge of the state of the atmosphere and ocean. It is hence possible that some of the historical sequences used might be inconsistent with current large-scale atmospheric conditions and would therefore be unlikely to occur in the next few months.

Period: January 2025 – March 2025


Issued on 07.01.2025 using data to the end of December 2024

The outlook for January indicates that flows are likely to be normal to below normal across southern and central England, except for a handful of slow-responding catchments which are likely to have normal or above normal flows. River flows in northern England and northern Scotland are likely to be normal to above normal. Elsewhere in the UK, flows are likely to be in the normal range. The January to March outlook indicates that flows across northern parts of the UK are likely to be normal to above normal, whilst southern areas are likely to experience flows within the normal range.


The historical weather analogues method uses Met Office predictions of average weather 1 and 3 months ahead to provide inputs to a hydrological model. Like the ESP method, observed rainfall and temperature data from past years are used to drive the predictions, however, the analogue method constrains the selection of past rainfall using the weather conditions in the meteorological forecasts (which are summarised for this forecast in the Met Office likelihood of impacts blocks underneath the maps). For each member of the Met Office forecast ensemble, the 10 analogues that best match the predicted average weather pattern (surface pressure map) over the forecast period are selected. Precipitation and temperature sequences constructed from the selected analogue scenarios are corrected to account for historic trends and used as inputs to hydrological models. Here, the GR6J model is run using these inputs, creating an ensemble of hydrological forecasts.

The outputs shown in the maps are the likelihoods of different outcomes for the average river flow over the one-month and three-month forecast periods at each location. The outlooks maps show the distribution for 64 catchments across the United Kingdom. Each bar plot represents the likelihood of the simulated river flow compared to the historical river flow, for the same n-month period. The probabilities fall within five categories, classified as: low, below normal, normal, above normal and high. The expected climatological probability of ensemble members in each of these categories is shown under the legend.

1-month river flow outlook starting Jan 2025



3-month river flow outlook starting Jan 2025



Met Office 1-month likelihood of precipitation impact



© Crown copyright, Met Office


Met Office 3-month likelihood of precipitation impact



© Crown copyright, Met Office

This page shows the **probability of extreme conditions in Scotland** over the next month. The bar chart indicates the likelihood of extremely high or low flows in a given number of regions. Maps show which regions are most likely to experience these flows as overall conditions get more extreme.

How likely are we to experience extremely low or high flows?



Probability of low and high flows in multiple regions simultaneously


0.0%

Summary

Scotland – one month


Forecasts suggest that river flows are most likely to be in the *normal* range for Scotland over the next month. River flows here are unlikely to be extremely high or low over the next three months. If extremely high flows are observed, they are most likely to occur in northwest Scotland

Which regions are most likely to experience extremely low or high flows?



This page shows the **probability of extreme conditions in Scotland** over the next three months. The bar chart indicates the likelihood of extremely high or low flows in a given number of regions. Maps show which regions are most likely to experience these flows as overall conditions get more extreme.

How likely are we to experience extremely low or high flows?



Probability of low and high flows in multiple regions simultaneously


0.7%

Summary

Scotland – three months

Forecasts suggest that river flows are most likely to be in the *normal* range for Scotland over the next month. Flows are unlikely to be *extremely low* over the next three months, however there is a good chance of *extremely high* flows in some Scottish regions.

Which regions are most likely to experience extremely low or high flows?




This page shows the **probability of extreme conditions in England and Wales** over the next month. The bar chart indicates the likelihood of extremely high or low flows in a given number of regions. Maps show which regions are most likely to experience these flows as overall conditions get more extreme.

Summary

England and Wales – one month

Forecasts suggest that river flows are most likely to be in the *normal* range for England and Wales over the next month. *Extremely high/low* river flows are less likely over the next month.


How likely are we to experience extremely low or high flows?



Probability of low and high flows in multiple regions simultaneously

0.0%

Which regions are most likely to experience extremely low or high flows?




This page shows the **probability of extreme conditions in England and Wales** over the next three months. The bar chart indicates the likelihood of extremely high or low flows in a given number of regions. Maps show which regions are most likely to experience these flows as overall conditions get more extreme.

Summary

England and Wales – three months

Forecasts suggest that river flows are most likely to be in the *normal* range for England and Wales over the next three months. If *extremely high flows* occur, they are most probable in northern England.


How likely are we to experience extremely low or high flows?



Probability of low and high flows in multiple regions simultaneously

1.0%

Which regions are most likely to experience extremely low or high flows?



P(High and low flows simultaneously) 1.0%

Histograms of GB regional mean river flow distributions


Period: January 2025 - March 2025

Issue date: 06.01.2025


- This page shows the **ensemble flow distribution for each region** over the next 1- and 3-month periods.
- The 1km flow forecasts are averaged within each region, then categorised according to the historical flows.
- The histogram in each region indicates the proportion of ensemble members in each flow category.

SUMMARY: Over the next month, river flows are likely to be in the *normal range*. River flows in the south are more likely to be in the *below normal to normal range*, while river flows in the north are more likely to be in the *normal to above normal range*.

Over the next three months, river flows in the north of Great Britain are likely to be in the *normal to exceptionally high range*. In Wales and southern and central England, river flows are likely to be in the *normal range*.




- SCOTLAND**
- HR Highlands Region
 - NER North East Region
 - TR Tay Region
 - FR Forth Region
 - CR Clyde Region
 - TWR Tweed Region
 - SR Solway Region
- ENGLAND**
- N Northumbria
 - NW North West
 - Y Yorkshire
 - ST Severn Trent
 - A Anglian
 - T Thames
 - S Southern
 - W Wessex
 - SW South West
- WALES**
- WEL Welsh



Flow estimate for each river pixel ranked in terms of historic % flow estimates (1963-2016)

Exceptionally high flow	> 95
Notably high flow	87-95
Above normal	72-87
Normal range	28-72
Below normal	13-28
Notably low flow	5-13
Exceptionally low flow	< 5




Tables of GB regional mean river flow distributions

Period: January 2025 - March 2025

Issue date: 06.01.2025

- This page shows **the ensemble flow distribution for each region** over the next 1- and 3-month periods.
- The 1km flow forecasts are averaged within each region, then categorised according to the historical flows.
- The table indicates the percentage of ensemble members in each flow category.



SUMMARY: Over the next month, river flows are likely to be in the *normal range*. River flows in the south are more likely to be in the *below normal to normal range*, while river flows in the north are more likely to be in the *normal to above normal range*.

Over the next three months, river flows in the north of Great Britain are likely to be in the *normal to exceptionally high range*. In Wales and southern and central England, river flows are likely to be in the *normal range*.


1-month ahead	A	NW	N	ST	SW	S	T	WEL	W	Y	CR	FR	HR	NER	SR	TR	TWR
Exceptionally high flow	3	12	10	6	2	1	6	4	6	5	9	10	16	10	3	3	7
Notably high	6	6	5	7	3	4	7	2	2	8	9	10	12	11	8	13	6
Above normal	9	21	9	13	6	12	9	12	7	18	10	12	13	15	21	12	11
Normal range	54	43	43	42	30	43	56	49	36	39	58	53	54	44	46	50	51
Below normal	24	10	24	20	42	39	23	20	44	22	10	9	5	14	12	8	19
Notably low	3	6	8	12	11	1	0	12	5	8	4	6	0	4	9	9	3
Exceptionally low flow	0	1	1	0	7	0	0	1	0	1	0	1	0	0	1	4	3

3-month ahead	A	NW	N	ST	SW	S	T	WEL	W	Y	CR	FR	HR	NER	SR	TR	TWR
Exceptionally high flow	0	27	19	12	1	3	3	14	2	13	15	23	22	9	17	8	26
Notably high	10	9	11	15	4	9	20	9	17	9	11	9	14	17	16	14	9
Above normal	21	12	17	14	13	19	14	13	10	27	28	20	18	22	22	13	10
Normal range	50	39	37	45	39	52	49	40	53	36	36	39	42	34	33	47	39
Below normal	18	12	5	9	22	16	13	16	11	10	6	4	4	12	8	9	11
Notably low	1	0	6	1	15	1	0	6	4	2	2	4	1	3	2	8	3
Exceptionally low flow	0	1	6	4	7	1	0	3	2	4	2	0	0	2	2	2	3

Forecasts of river flows using Met Office rainfall forecasts

- The data on these (yellow-bordered) pages are based on approximately 400 rainfall scenarios provided by the Met Office, which are used as inputs to a water balance hydrological model.
- River flow forecasts for every 1km grid cell are ranked according to the historical flow estimates and aggregated within each region.
- A full description of this method and these summary products is given in the technical documentation available via the Hydrological Outlook website.

Probability of extreme river flows





Extreme river flows are defined as those which rank in the lowest or highest 13% of historical flow estimates (1963 - 2016). This definition encompasses the 'Notably' and 'Exceptionally' high/low flow bands used elsewhere in the Outlook.

The bar chart shows the probability of a given number of regions experiencing extremely high/low flows, where scenarios showing both extremely high and extremely low flows in different regions simultaneously excluded. These probabilities are also shown beneath the maps.

Shaded regions on each map are those most likely to experience extreme flows from the set of scenarios with at least a given number of regions experiencing such flows. If shown, grey maps indicate scenarios not observed in the ensemble.

The box drawn around some maps spans the central 50% probability interval, excluding scenarios where extremely high/low flows are observed simultaneously. If these excluded cases constitute a significant probability, details are given in the yellow summary box.

Regional mean river flow distributions



The maps illustrate the ensemble distribution of regional mean river flows. The historical distribution is shown at bottom-left, and allows deviations from the normal distribution to be determined by comparing the forecast distribution to the historical distribution. A summary is given in the yellow box.

The table gives access to the data shown in the histograms. The numbers in the tables are the percentage of ensemble forecasts falling in each of the flow categories. As before results are averaged by region then ranked in terms of 54 years of historical regional flow estimates (1963 – 2016).


Period: January 2025 – March 2025

Issued on 08.01.2025 using data to the end of November.


Under median rainfall conditions, groundwater levels are expected to remain above normal to exceptionally high across most aquifers over the next month, including the Yorkshire Chalk, Jurassic Limestones, and the Chilterns. Normal levels are anticipated at some sites in southern England and Wales during this period. The three-month median forecasts show more variation, with groundwater levels becoming more normal in East Anglia, Lincolnshire, and Central England. However, in parts of the Wessex Chalk and South Downs Chalk, levels are expected to rise to above normal or notably high.

These forecasts are produced by running five members of the Met Office ensemble climate forecast through groundwater models of observation borehole hydrographs at 42 sites across the country. The sites are distributed across the principal aquifers.

Based on the distribution of observed historical groundwater levels in a given month, seven categories have been derived for each site: very low, low, below normal, normal, above normal, high, and very high. The forecast groundwater level is assigned to one of these seven categories depending on where it falls within the distribution of the historically observed values.




1-month outlook



3-month outlook

Key	Percentile range of historic observed values for relevant month
Exceptionally high levels	> 95
Notably high levels	87-95
Above normal	72-87
Normal	28-72
Below normal	12-28
Notably low levels	5-13
Exceptionally low levels	< 5






Outlook based on modelled groundwater from historical climate

Period: January 2025 – December 2025

Issued on 08.01.2025 using data to the end of November.

Groundwater levels at Skirwith in the Permo-Triassic Sandstones are forecast to remain at above normal to exceptionally high levels for the next nine months, followed by a transition to more normal levels. In the Chalk at Rockley, the Jurassic Limestone at New Red Lion and the Permo-Triassic Sandstones at Bussels No 7A, groundwater levels are likely to transition towards normal conditions over the next three to six months. In the Chalk at Little Bucket Farm and Dalton Holme, groundwater levels are likely to remain normal over the next 12 months, but with a possibility of both sites experiencing periods of above or below normal conditions at times during this period.



This outlook is based on monthly ensembles of historical sequences of observed climate (rainfall and potential evapotranspiration) that form input to hydrological models. The outputs are probabilistic simulations of the average groundwater level over the forecast horizon (3 to 12 months ahead), at each location.

that fall within each the seven categories: exceptionally low, notably low, below normal, normal, above normal, notably high and exceptionally high. The monthly variations can be compared to the long-term average distribution of levels, which are shown as columns on the left and right of each graph.

the atmosphere and ocean. It is hence possible that some of the historical sequences used might be inconsistent with current large-scale atmospheric conditions and would therefore be unlikely to occur in the next few months.

The graphs show variation over time of the number of simulated groundwater levels in each monthly ensemble,

This outlook is based entirely on historical sequences and therefore does not contain any knowledge of the state of