


This map shows the simulated monthly mean flow across Great Britain for last month, ranked in terms of 54 years of historical flow estimates (1963 – 2016).

These flows are produced by the 1km resolution Grid-to-Grid (G2G) hydrological model, which is run up to the end of each calendar month using observed rainfall and MORECS potential evaporation as input.

Note that the G2G model provides estimates of natural flows.



Issue date: 03.02.2026

February 2026

These maps are based on Grid-to-Grid (G2G) hydrological model simulated subsurface water storage (water in the soil and groundwater), expressed as an anomaly from the historical monthly mean. To highlight areas that are particularly wet or dry, the storage anomaly is presented relative to historical extremes. Rainfall in WET areas with high positive relative wetness could result in flooding in the coming days/weeks. Areas of negative relative wetness indicate locations which are particularly DRY, and little or no rain in these areas could potentially lead to (or prolong) a drought. Maps of soil moisture only are available on the next page.

**SUMMARY:** Subsurface water stores are very high (wet) across much of England and eastern Scotland, but lower than usual in western Scotland and the north west of England.




## Relative wetness

Water storage anomaly as a % of maximum (positive wetness) or minimum (negative wetness) storage anomaly (zero indicates average value)

High (very wet)	>80
	60-80
Medium wetness	40-60
	20-40
Low wetness	0.01-20
Average conditions	-0.01-0.01
Low dryness	-20-0.01
	-40-20
Medium dryness	-60-40
	-80-60
High (very dry)	<-80

Major rivers

Labels refer to estimated storage on final day of named month




\*Example month displaying extreme positive wetness

February 2026

These maps are based on Grid-to-Grid (G2G) hydrological model simulated soil moisture, expressed as an anomaly from the historical monthly mean. To highlight areas that are particularly wet or dry, the soil moisture anomaly is presented relative to historical extremes. These maps are not a forecast; rather an indication of current conditions. Soil moisture will often look similar to total storage (shown on the previous slide), since total storage comprises both soil moisture and storage in the saturated zone.

**SUMMARY:** Soil water stores are very high (wet) across much of England and eastern Scotland, but lower than usual in western Scotland and the north west of England.






These maps show the Grid-to-Grid (G2G) hydrological model simulated subsurface water storage, expressed as an anomaly from the historical monthly mean (1981-2010), presented on a 1km grid and as regional means. Subsurface storage deficits, i.e. where the subsurface water storage anomaly is less than zero, are highlighted in red/pink.

The subsurface storage deficit (mm) can be interpreted as an estimate of additional rainfall that would be required in future months to overcome dry conditions (i.e. rainfall in addition to what is expected on average). Regional mean values of additional rainfall required are provided in the table below.


## Regional estimate of additional rainfall required (mm)

### SCOTLAND

26	HR	Highlands Region
0	NER	North East Region
0	TR	Tay Region
6	FR	Forth Region
9	CR	Clyde Region
0	TWR	Tweed Region
0	SR	Solway Region



Water storage deficit (anomaly; mm)




These maps show the return period of the rainfall required to overcome dry conditions simulated using the Grid-to-Grid (G2G) hydrological model. The maps are coloured according to the return period of accumulated rainfall required to overcome the estimated current subsurface water storage deficit over the next one to six months (areas with no storage deficit will always be white). These maps do not provide a drought forecast; instead they indicate whether particularly heavy rainfall would be required to return to normal conditions for the time of year.

## SUMMARY:

Subsurface deficits in the Highlands are most likely to return to normal within the next month, but if they do not are unlikely to recover for a few months.

The small deficits present in other regions do not require unusually high rainfall to recover to normal for this time of year.



## SCOTLAND

HR Highlands Region

NER North East Region

TR Tay Region

FR Forth Region

CR Clyde Region

TWR Tweed Region

SR Solway Region

## ENGLAND

N Northumbria

NW North West

Y Yorkshire

ST Severn Trent

A Anglian

T Thames

S Southern

W Wessex

SW South West

## WALES

WEL Welsh



February 2026

CURRENT CONDITIONS