

Hydrological Outlook UK

Period: From April 2020

Issued on 08.04.2020 using data to the end of March 2020

SUMMARY

The outlook is for river flows in northern and western parts of the UK to be normal to below normal for April and over the next three months. Flows across central southern England and the East Midlands are likely to be normal to above normal for April, and within the normal range for the three month period April-June. Both river flows and groundwater levels are expected to be within the normal range in East Anglia for April and over the three months to June. Elsewhere, groundwater levels are likely to be normal to above normal for April, and normal over the next three months, with a few localised exceptionally high levels expected in the Permo-Triassic sandstones.

Rainfall:

Rainfall in March was a stark contrast to the record-breaking wet February, with drier than average weather seen across much of the UK. The vast majority of the UK saw less than 90% of average rainfall, whilst much of the eastern UK saw less than 50% of average.

The rainfall outlook (issued by the Met Office on 26th March 2020) is that for April, the likelihoods of above- and below-average precipitation are similar. For April-May-June as a whole, below-average precipitation is moderately more likely than above-average precipitation. The probability that UK-average precipitation for April-May-June will fall into the driest of five equal categories is 25% and the probability that it will fall into the wettest of the five categories is around 15% (the 1981-2010 probability for each of the categories is 20%).

River flows:

River flows in March generally dropped from the exceptionally high levels seen in February and though reduced, flows still remained above normal to exceptionally high across much of the UK. Flows in East Anglia and most of Scotland were generally within the normal range for March, and a few catchments in north-eastern Scotland and Northern Ireland were below normal.


Following the dry weather in March, that is likely to continue over the next three months, the outlook is for flows to continue to drop across the northern and western parts of the UK. Flows in this region are therefore likely to be normal to below normal over for April and over the next three months to June. River flows across central southern England and the East Midlands are likely to remain normal to above normal for April, but river flows in this area, as well as East Anglia, are likely to be within the normal range for April-May-June as a whole.

Groundwater:

As with river flows, groundwater levels dropped in many boreholes across the UK in March. However several aquifers still recorded exceptionally high and record breaking high levels, in particular in the Permo-Triassic sandstones of central and northern England, and the Chalk aquifers of southern England. Groundwater levels in East Anglia were normal to below normal.

The outlook is for groundwater levels across the UK to continue recessing in most areas, to be normal to above normal for April, though levels in East Anglia are most likely to be within their normal range. Over the three month period April-June, levels are generally expected to return to being within their normal range across the UK, however some localised aquifers are expected to remain exceptionally high, particularly in the Permo-Triassic sandstones.

The Hydrological Outlook UK provides an outlook for the water situation for the UK over the next three months and beyond. For guidance on how to interpret the outlook, a wider range of information, and a full description of underpinning methods, please visit the website: www.hydoutuk.net



Hydrological Outlook UK

About the Hydrological Outlook:

This document presents an outlook for the UK water situation for the next 1 – 3 months and beyond, using observational datasets, meteorological forecasts and a suite of hydrological modelling tools. The outlook is produced in a collaboration between the UK Centre for Ecology and Hydrology (UKCEH), British Geological Survey (BGS), the Met Office, the Environment Agency (EA), Natural Resources Wales (NRW), the Scottish Environment Protection Agency (SEPA), and for Northern Ireland, the Department for Infrastructure – Rivers (DfIR).

Data and Models:

The Hydrological Outlook depends on the active cooperation of many data suppliers. This cooperation is gratefully acknowledged. Historic river flow and groundwater data are sourced from the UK National River Flow Archive and the National Groundwater Level Archive. Contemporary data are provided by the EA, SEPA, NRW and DfIR. These data are used to initialise hydrological models, and to provide outlook information based on statistical analysis of historical analogues.

Climate forecasts are produced by the Met Office. Hydrological modelling is undertaken by UKCEH using the Grid-to-Grid, PDM and CLASSIC hydrological models and by the EA using CATCHMOD. Hydrogeological modelling uses the R-groundwater model run by BGS and CATCHMOD run by the EA. Supporting documentation is available from the Outlooks website:

<http://www.hydoutuk.net/methods>

Presentation:

The language used in the summary presented overleaf generally places flows and groundwater levels into just three classes, i.e. below normal, normal, and above normal. However, the underpinning methods use as many as seven classes as defined in the graphic to the right, i.e. the summary uses a simpler classification than some of the methods. On those occasions when it is appropriate to provide greater discrimination at the extremes the terminology and definitions of the seven class scheme will be adopted.

Percentile range of historic values for relevant month

Exceptionally high flow	> 95
Notably high flow	87-95
Above normal	72-87
Normal range	28-72
Below normal	13-28
Notably low flow	5-13
Exceptionally low flow	< 5

Disclaimer and liability:

The Hydrological Outlook partnership aims to ensure that all Content provided is accurate and consistent with its current scientific understanding. However, the science which underlies hydrological and hydrogeological forecasts and climate projections is constantly evolving. Therefore any element of the Content which involves a forecast or a prediction should not be relied upon as though it were a statement of fact. To the fullest extent permitted by applicable law, the Hydrological Outlook Partnership excludes all warranties or representations (express or implied) in respect of the Content.

Your use of the Content is entirely at your own risk. We make no warranty, representation or guarantee that the Content is error free or fit for your intended use.

From April 2018 the Hydrological Outlook is supported by the Natural Environment Research Council funded [UK-SCAPE](#) and [Hydro-JULES](#) Programmes.

Copyright:

Some of the features displayed on the maps contained in this report are based on the following data with permission of the controller of HMSO.

- (i) Ordnance Survey data. © Crown copyright and/or database right 2005. Licence no. 100017897.
- (ii) Land and Property Services data. © Crown copyright and database right, S&LA 145.
- (iii) Met Office rainfall data. © Crown copyright.

All rights reserved. Unauthorised reproduction infringes crown copyright and may lead to prosecution or civil proceedings.

Further information:

For more detailed information about the Hydrological Outlook, and the derivation of the maps, plots and interpretation provided in this outlook, please visit the Hydrological Outlook UK website.

The website features a host of other background information, including a wider range of sources of information which are used in the preparation of this Outlook.

Contact:

Hydrological Outlooks UK, UK Centre for Ecology & Hydrology, Wallingford, Oxfordshire, OX10 8BB
t: 01491 692371 e: enquiries@hydoutuk.net

Reference for the Hydrological Outlook:

Hydrological Outlook UK, 2020, April, UK Centre for Ecology and Hydrology, Oxfordshire UK, Online, <http://www.hydoutuk.net/latest-outlook/>

Other Sources of Information:

The Hydrological Outlook should be used alongside other sources of up-to-date information on the current water resources status and flood risk.

Environment Agency Water Situation Reports: provides summary of water resources status on a monthly and weekly basis for England:

<https://www.gov.uk/government/collections/water-situation-reports-for-england>

Flood warnings are continually updated, and should be consulted for an up-to-date and localised assessment of flood risk:

Environment Agency: <https://flood-warning-information.service.gov.uk/map>

Scottish Environment Protection Agency: <http://www.sepa.org.uk/flooding.aspx>

Hydrological Summary for the UK: provides summary of current water resources status for the UK:

<https://nrfa.ceh.ac.uk/monthly-hydrological-summary-uk>

UK Met Office forecasts for the UK:

www.metoffice.gov.uk/public/weather/forecast/#?tab=regionalForecast

UK Water Resources Portal: monitor the UK hydrological situation in near real-time including rainfall, river flow, groundwater and soil moisture from COSMOS-UK:

<https://eip.ceh.ac.uk/hydrology/water-resources/>



Met Office 3-month Outlook

Period: April – June 2020 Issue date: 26.03.20

The forecast presented here is for April and the average of the April-May-June period for the United Kingdom as a whole. The forecast for April will be superseded by the long-range information on the public weather forecast web page (www.metoffice.gov.uk/public/weather/forecast/#?tab=regionalForecast), starting from 1st April 2020. This forecast is based on information from observations, several numerical prediction systems and expert judgement.

SUMMARY – PRECIPITATION:

For April, the likelihoods of above- and below-average precipitation are similar. For April-May-June as a whole, below-average precipitation is moderately more likely than above-average precipitation.

The probability that UK-average precipitation for April-May-June will fall into the driest of our five categories is 25% and the probability that it will fall into the wettest of our five categories is around 15% (the 1981-2010 probability for each of these categories is 20%).

CONTEXT:

There are fewer global drivers of UK weather patterns at this time of year compared to the winter season, so predictability of precipitation amounts is lower. This means there are typically only small shifts in the likelihood of above- and below-average precipitation. As discussed in the Outlook for temperature, there is a greater-than-usual likelihood of a positive phase of the North Atlantic Oscillation (NAO), particularly early in the Outlook period. This would favour above-average precipitation. Nevertheless, there are also signals for high pressure near the UK, which favour drier-than-average conditions.

For April, prediction systems show that the chances of above- and below-average rainfall are balanced, indicating the uncertainty over which of the factors above will exert more influence (see left-hand graph of figure P2). For the April-May-June period, below-average precipitation is moderately more likely than above-average precipitation, consistent with a greater likelihood that high-pressure will be more influential during the Outlook period overall (see right-hand graph of figure P2). This does not preclude spells of wet and windy weather occurring at times during the period.

Fig P2

1-month and 3-month UK outlook for precipitation in the context of observed climatology




Fig P1

3-month UK outlook for precipitation in the context of the observed annual cycle





Fig P3

1-month and 3-month UK outlook for precipitation in the context of recent climatology: year-to-year and within-season variability



This Outlook provides an indication of possible temperature and rainfall conditions over the next 3 months. It is part of a suite of forecasts designed for contingency planners.

The Outlook should not be used in isolation but should be used with shorter-range and more detailed (30-day, 15-day and 1-to-7-day) forecasts and warnings available to the contingency planning community from the Met Office.



Met Office 3-month Outlook

Period: April – June 2020 Issue date: 26.03.20

The forecast presented here is for April and the average of the April-May-June period for the United Kingdom as a whole. The forecast for April will be superseded by the long-range information on the public weather forecast web page (www.metoffice.gov.uk/public/weather/forecast/#?tab=regionalForecast), starting from 1st April 2020. This forecast is based on information from observations, several numerical prediction systems and expert judgement.

SUMMARY – TEMPERATURE:

For April, above-average temperatures are slightly more likely than below-average temperatures. For April-May-June as a whole, above-average temperatures are more likely than below-average temperatures.

Overall, the probability that the UK-average temperature for April-May-June will fall into the coldest of our five categories is less than 5% and the probability that it will fall into the warmest of our five categories is between 40 and 45% (the 1981-2010 probability for each of these categories is 20%).

CONTEXT:

The El Niño-Southern Oscillation (ENSO) is currently in a neutral phase, with little likelihood of a significant El Niño or La Niña event developing during the Outlook period. Consequently, it is not expected to influence UK weather patterns.

The Stratospheric Polar Vortex (SPV) – the circulation of winds in the stratosphere above the Arctic – has recently been very strong and is set to remain stronger than usual in the remaining weeks before its seasonal breakdown in April. This, along with the lag of about 2 weeks in responses in the lower layers of the atmosphere, favours a positive phase of the North Atlantic Oscillation (NAO) during the first half of the Outlook period.

Whilst a positive phase of the NAO favours milder-than-normal conditions, its effect on temperatures is less marked during the next 3-months than in winter. Sea surface temperatures to the west of the UK are additionally below average, further moderating the warm signal.

For April and April-May-June as a whole, the Met Office long-range prediction system has good support from predictions from other centres around the world, showing an increased likelihood that the UK will come under the influence of high pressure as well as westerly winds at times. High pressure favours above-average temperatures during late spring and early summer, and with the warming of climate, this results in an increase in the probability of above-average temperatures (see graphs of figure T2).

The relatively high probability of our warmest forecast category does not imply extreme or unseasonal weather throughout the whole 3-month period. Indeed, the Outlook does not identify weather for a particular day or week. The increased likelihood of this category could mean more days with temperatures that are above average to a more modest degree. Above-average temperatures can also arise from a range of types of weather, not just sunny and dry conditions.

Fig T1




Fig T2

1-month and 3-month UK outlook for temperature in the context of observed climatology





Fig T3

1-month and 3-month UK outlook for temperature in the context of recent climatology: year-to-year and within-season variability




This Outlook provides an indication of possible temperature and rainfall conditions over the next 3 months. It is part of a suite of forecasts designed for contingency planners.

The Outlook should not be used in isolation but should be used with shorter-range and more detailed (30-day, 15-day and 1-to-7-day) forecasts and warnings available to the contingency planning community from the Met Office.

SUMMARY

The outlooks for April and for April-June are for normal to above normal river flows across most of the UK, with the exception of north-east Scotland where normal to below normal flows are likely. Please note that not many forecasts are available for the west of the country.

River flow outlook for Apr 2020




1-month flow outlook

Outlooks from hydrological analogues are based on a comparison of river flow during recent months with flows during the same months in previous years at a set of approximately 90 sites from across the UK. These sites are depicted on the two maps. Years with observed flows that most closely resemble current conditions are identified as the best analogues and the outlook is based on extrapolating from current conditions based on these analogues.

It is, however, often the case that a simpler forecast based on the persistence of river flow provides a better forecast than provided by analogy. This is particularly true for slowly responding catchments associated with aquifer outcrops.

Both methods are considered at each site and the forecast from the method with the higher confidence is presented. A simple classification of flows is used (high, medium and low) as indicated by the colours of the dots, with the confidence

River flow outlook for Apr - Jun 2020



3-month flow outlook

of the forecast being represented by the size of the dot. A tag on the dot indicates which method has been used in each instance.




Period: April 2020

Issued on 06.04.2020 using data to the end of March 2020

These figures provide insight into the hydrological analogue methodology for a set of sites from across the UK.

In each of the time series graphs the bold black line represents the observed flow during the past six months. The grey band indicates the normal flow range (the normal band includes 44%

of observed flows in each month). The selected analogues are shown as thin lines and the trajectories that flows took in the following month are also shown. The forecast is shown as the dashed red line, and in each plot it states whether this has come from the analogues or has been generated on the basis of persistence.











Period: April – June 2020

Issued on 06.04.2020 using data to the end of March 2020

These figures provide insight into the hydrological analogue methodology for a set of sites from across the UK.

In each of the time series graphs the bold black line represents the observed flow during the past nine months. The grey band indicates the normal flow range (the normal band includes 44%

of observed flows in each month). The selected analogues are shown as thin lines and the trajectories that flows took in the following three months are also shown. The forecast is shown as the dashed red line, and in each plot it states whether this has come from the analogues or has been generated on the basis of persistence.




Period: April 2020 – June 2020

Issued on 07.04.2020 using data to the end of March


Normal to above normal groundwater levels are predicted to prevail over the UK in the next month. Exceptionally high groundwater levels are predicted in some sites in the Chalk of the South of England and in the Permo-Triassic sandstones of the North-West. In the 3 month forecast, more normal conditions are predicted throughout majority of the UK, with few individual sites predicting below normal or exceptionally high levels. Above normal conditions are predicted to prevail through the Chalk of the North Downs and some areas of Chalk in the South of England. Note there are a reduced number of modelled sites. This is due to the temporary unavailability of data, where EA staff have been unable to either manually dip boreholes or download logger data as a consequence of Covid-19 restrictions.

These forecasts are produced by running five members of the Met Office ensemble climate forecast through groundwater models of observation borehole hydrographs at 42 sites across the country. The sites are distributed across the principal aquifers.

Based on the distribution of observed historical groundwater levels in a given month, seven categories have been derived for each site: very low, low, below normal, normal, above normal, high, and very high. The forecast groundwater level is assigned to one of these seven categories depending on where it falls within the distribution of the historically observed values.



Key	Percentile range of historic observed values for relevant month
Exceptionally high levels	> 95
Notably high levels	87-95
Above normal	72-87
Normal	28-72
Below normal	13-28
Notably low levels	5-13
Exceptionally low levels	< 5



Current Daily Simulated Subsurface Water Storage Conditions

Based on subsurface water storage estimated for 31st March 2020


Issue date: 03.04.2020

These maps are based on Grid-to-Grid (G2G) hydrological model simulated subsurface water storage, expressed as an anomaly from the historical monthly mean. To highlight areas that are particularly wet or dry, the storage anomaly is presented here using a colour scale highlighting water storage relative to historical extremes. The maps below show relative dryness.

These maps do not provide a drought forecast and are not maps of soil moisture. Instead they indicate areas where subsurface water storage approaches or exceeds its historical minimum. A lack of rainfall in the high 'relative dryness' areas could lead to (or prolong) a drought.

SUMMARY: At the end of March, much of the country is experiencing relative dryness levels that are slightly higher than average for this time of year, with some small, scattered areas of high relative dryness.

Relative Dryness



The Hydrological Outlook UK provides an outlook for the water situation for the UK over the next three months and beyond. For guidance on how to interpret the outlook, a wider range of information, and a full description of underpinning methods, please visit the website: www.hydoutuk.net

Current Daily Simulated Subsurface Water Storage Conditions

Based on subsurface water storage estimated for 31st March 2020




Issue date: 03.04.2020

These maps are based on Grid-to-Grid (G2G) hydrological model simulated subsurface water storage, expressed as an anomaly from the historical monthly mean. To highlight areas that are particularly wet or dry, the storage anomaly is presented here using a colour scale highlighting water storage relative to historical extremes. The maps below show relative wetness.

These maps do not provide a flood forecast and are not maps of soil moisture. Instead they indicate areas where subsurface water storage approaches or exceeds its historical maximum. Rainfall in the high 'relative wetness' areas could result in flooding.

SUMMARY: At the end of March, while most of the country is not experiencing relative wetness levels above average for this time of year, some areas through central and southern England are still experiencing higher than average relative wetness.

Relative Wetness



*Example month displaying extreme relative wetness

Relative Dryness

- The relative dryness map highlights areas where current estimates of **subsurface water storage** (from the G2G hydrological model, calculated for the last day of last month) are particularly **low**.
- The map indicates areas where the ground is dry compared to the monthly **average** storage (for the period 1981 to 2010), and shows this relative to the historical **minimum** storage level (for 1971 to 2010).
- Relative dryness calculation:
$$R_d (\%) = \frac{(S_{average} - S)}{(S_{average} - S_{min})} \times 100$$

$$= \frac{(\text{average storage for this month} - \text{storage at end of last month})}{(\text{average storage for this month} - \text{historical minimum storage})} \times 100$$
- A value of $R_d = 100$ shows that a region is very dry, and indicates that the storage is as low as the minimum value ever estimated by the model for this month.
- A value of $R_d = 0$ indicates that the storage in the region matches the monthly average value. *Negative relative dryness values will show up as part of the relative wetness map.*
- The map **does not provide a drought forecast**. A lack of rainfall in the high 'relative dryness' areas **could** lead to (or prolong) a drought.

Relative Wetness

- The relative wetness map highlights areas where current estimates of **subsurface water storage** (from the G2G hydrological model, calculated for the last day of last month) are particularly **high**.
- The map indicates areas where the ground is wet compared to the monthly **average** storage (for the period 1981 to 2010), and shows this relative to the historical **maximum** storage level (for 1971 to 2010).
- Relative wetness calculation:
$$R_w (\%) = \frac{(S - S_{average})}{(S_{max} - S_{average})} \times 100$$








$$= \frac{(\text{storage at end of last month} - \text{average storage for this month})}{(\text{historical maximum storage} - \text{average storage for this month})} \times 100$$
- A value of $R_w = 100$ shows that a region is very wet, and indicates that the storage is as high as the maximum value ever estimated by the model for this month.
- A value of $R_w = 0$ indicates that the storage in the region matches the monthly average value. *Negative relative wetness values will show up as part of the relative dryness map.*
- The map **does not provide a flood forecast**. Rainfall in the high 'relative wetness' areas **could** result in flooding.

These maps show the **return period** of the rainfall required to overcome dry conditions simulated using the Grid-to-Grid (G2G) hydrological model. The maps are coloured according to the return period of accumulated rainfall required to overcome the estimated current subsurface water storage deficit over the next few months.

These maps do not provide a drought forecast. Instead they indicate the return period of rainfall required to overcome the dry conditions for the following 6 months based on current conditions.

SUMMARY: During April to September, Britain will not require particularly unusual rainfall (0 to 5 year return periods) to return to average conditions for the time of year.



Rainfall amount / Probability		Return period (years)	
Low (this rain is likely to occur)	> 20%		0 - 5
	< 20%		5 - 10
	< 10%		10 - 25
	< 4%		25 - 50
High (less likely)	< 2%		50 - 100
	< 1%		100 - 200
Extreme (unlikely but still possible)	< 0.5%		>200

SCOTLAND


HR Highlands Region
NER North East Region
TR Tay Region
FR Forth Region
CR Clyde Region
TWR Tweed Region
SR Solway Region

ENGLAND

N Northumbria
NW North West
Y Yorkshire
ST Severn Trent
A Anglian
T Thames
S Southern
W Wessex
SW South West

WALES

WEL Welsh



NORTHERN IRELAND

This method cannot currently be used in Northern Ireland

Method

- These maps show the **return period** of the rainfall required to overcome dry conditions simulated using the Grid-to-Grid (G2G) hydrological model. The maps are coloured according to the return period of rainfall required to overcome the estimated current subsurface water storage deficit.
- For **dry areas** within a Hydrological Outlook region, i.e. where subsurface water storage anomaly < 0 , we estimate **regional average subsurface water storage deficit** (mm) from the last day of the most recent G2G model run.
- For each region we also estimate the **regional monthly average rainfall total** (mm) (for the period 1971-2000).
- For each of the next 6 months, we estimate the rainfall total (including what is normally expected for each month) required to overcome the dry conditions.
 - To overcome the dry conditions by the end of month 1:
rainfall required (mm) = regional monthly average rainfall for month 1 + regional average storage deficit
 - To overcome the dry conditions by the end of month 2 (more likely):
rainfall required (mm) = regional monthly average rainfall for months 1 and 2 + regional average storage deficit
 - To overcome the dry conditions by the end of month n (likely):
rainfall required (mm) = regional monthly average rainfall for months 1 to n + regional average storage deficit
- Using Tabony tables we estimate the return period of the **rainfall required** in each region and over the next 1 to 6 months to overcome the dry conditions.
- The return period results are displayed as regional maps with the colour scale based on the return period (years) of the rainfall required to replenish subsurface stores over the next 1, 2, ..., 6 months ahead.
- Note: These maps do not provide a drought forecast. Instead they indicate the return period of rainfall required to overcome the dry conditions for the following 6 months based on current conditions.

Estimate of Additional Rainfall Required to Overcome Dry Conditions

Based on subsurface water storage estimated for 31st March 2020

Issue date: 03.04.2020

These maps show the Grid-to-Grid (G2G) hydrological model simulated subsurface water storage, expressed as an anomaly from the historical monthly mean (1981-2010), presented on a 1km grid and as regional means.

Subsurface storage deficits, i.e. where the subsurface water storage anomaly is less than zero, are highlighted by the red/pink colours.

The **subsurface storage deficit (mm)** can be interpreted as an estimate of additional rainfall that would be required in future months to overcome dry conditions (i.e. rainfall in addition to what is expected on average). Regional mean values of additional rainfall required are provided in the table below.

Regional estimate of additional
rainfall required (mm)

SCOTLAND




8	HR	Highlands Region
19	NER	North East Region
6	TR	Tay Region
6	FR	Forth Region
14	CR	Clyde Region
7	TWR	Tweed Region
17	SR	Solway Region

ENGLAND


9	N	Northumbria
13	NW	North West
1	Y	Yorkshire
0	ST	Severn Trent
0	A	Anglian
0	T	Thames
0	W	Wessex
0	S	Southern
16	SW	South West

WALES

11	WEL	Welsh
----	-----	-------



Water storage deficit
(anomaly, mm)



Period: April 2020 – June 2020

Issued on 03.04.2020 using data to the end of March

SUMMARY: During April, river flows across England and Wales are most likely to be in the *Normal range*. River flows in Scotland are most likely to be in the *Normal range* or below.

Over the next 3 months river flows across the country are most likely to be in the *Normal range* or below.

These forecasts are produced by using five members of the Met Office rainfall forecast ensemble as input to a water balance hydrological model to provide the five estimates of river flows shown on the left for one month and three months ahead.

Regional forecast monthly-mean river flows are derived from the average of 1km river flow estimates within each region and ranked in terms of 54 years of historical flow estimates (1963 – 2016).

The five maps illustrate the wide range of possible flows and while there is a 50% chance of flows between the 1st and 3rd quartiles, actual flows may be more extreme than the flows derived using the highest or lowest rainfall forecasts.

1-month flow outlook



Lowest rainfall forecast

1st quartile

Median

3rd quartile

Highest rainfall forecast



Key

Exceptionally high flow
Notably high flow
Above normal
Normal range
Below normal
Notably low flow
Exceptionally low flow

Percentile range of historic values for relevant month

> 95
87-95
72-87
28-72
13-28
5-13
< 5

3-month flow outlook





Lowest rainfall forecast

1st quartile

Median

3rd quartile

Highest rainfall forecast



SCOTLAND

HR Highlands Region
NER North East Region
TR Tay Region
FR Forth Region
CR Clyde Region
TWR Tweed Region
SR Solway Region

ENGLAND

N Northumbria
NW North West
Y Yorkshire
ST Severn Trent
A Anglian
T Thames
S Southern
W Wessex
SW South West

WALES

WEL Welsh



NORTHERN IRELAND

This method cannot currently be used in Northern Ireland

Period: April 2020 – June 2020


Issue date: 03.04.2020

The regional maps illustrating the regional river flows for five members of the Met Office ensemble of rainfall forecasts give some indication of the range of possible river flows in the coming months. As noted previously, the actual flows could be more extreme than the flows generated by either the lowest or highest members of the rainfall ensemble.

The bar charts (below) give further insight into the range of river flow forecasts by considering all members of the forecast rainfall ensemble. The regional bar charts show the percentage of ensemble forecasts falling in each of the flow categories as generated by the monthly-resolution water-balance model. As before results are averaged by region then ranked in terms of 54 years of historical regional flow estimates (1963 – 2016).

SUMMARY: During April, river flows across England and Wales are most likely to be in the *Normal range*. River flows in Scotland are most likely to be in the *Normal range* or below.

Over the next 3 months river flows across the country are most likely to be in the *Normal range* or below.



Period: April 2020 – June 2020

Issue date: 03.04.2020

The maps illustrating the regional river flows for five members of the Met Office ensemble of rainfall forecasts give some indication of the range of possible river flows in the coming months. As noted previously, the actual flows could be more extreme than the flows generated by either the lowest or highest members of the rainfall ensemble.

The tables below give further insight into the range of river flow forecasts by considering all members of the forecast rainfall ensemble. The numbers in the tables are the percentage of ensemble forecasts falling in each of the flow categories as generated by the monthly-resolution water-balance model. As before results are averaged by region then ranked in terms of 54 years of historical regional flow estimates (1963 – 2016).

SUMMARY: During April, river flows across England and Wales are most likely to be in the *Normal range*. River flows in Scotland are most likely to be in the *Normal range* or below.

Over the next 3 months river flows across the country are most likely to be in the *Normal range* or below.

SCOTLAND

HR Highlands Region
NER North East Region
TR Tay Region
FR Forth Region
CR Clyde Region
TWR Tweed Region
SR Solway Region

ENGLAND

N Northumbria
NW North West
Y Yorkshire
ST Severn Trent
A Anglian
T Thames
S Southern
W Wessex
SW South West

WALES

WEL Welsh



NORTHERN IRELAND

This method cannot currently be used in Northern Ireland

1-month ahead	A	NW	N	ST	SW	S	T	Welsh	W	Y	CR	FR	HR	NER	SR	TR	TWR
Exceptionally high flow	0	2	2	2	0	0	0	2	2	0	2	2	17	0	2	2	2
Notably high flow	2	2	0	7	2	2	5	0	0	2	14	2	0	2	12	2	0
Above normal	14	12	19	17	14	12	14	17	33	14	5	14	5	7	7	12	14
Normal range	79	57	60	71	67	69	81	64	62	69	38	55	52	60	48	52	69
Below normal	5	12	19	2	17	17	0	17	2	14	26	14	12	14	17	19	14
Notably low flow	0	14	0	0	0	0	0	0	0	0	14	12	14	12	14	12	0
Exceptionally low flow	0	0	0	0	0	0	0	0	0	0	0	0	0	5	0	0	0


3-months ahead	A	NW	N	ST	SW	S	T	Welsh	W	Y	CR	FR	HR	NER	SR	TR	TWR
Exceptionally high flow	0	5	0	0	0	0	0	5	0	0	2	5	5	0	0	0	0
Notably high flow	0	0	5	5	2	2	2	0	7	0	2	0	2	2	5	5	5
Above normal	10	5	5	7	7	2	7	5	2	10	5	5	5	5	7	5	5
Normal range	74	33	55	57	55	81	88	45	86	50	40	33	45	26	36	31	55
Below normal	17	29	26	31	26	14	2	33	5	38	29	48	29	33	29	45	14
Notably low flow	0	29	10	0	10	0	0	10	0	2	21	10	12	21	24	12	21
Exceptionally low flow	0	0	0	0	0	0	0	2	0	0	0	0	2	12	0	2	0

Period: April 2020 – September 2020

Issued on 03.04.2020 using data to the end of March


River flows across the UK are expected to be within the normal range for the next one to three months. The exception to this is some slowly responding groundwater-fed catchments in central southern England, where flows are expected to remain above normal for the next three months.

1-month river flow outlook starting Apr 2020




This outlook is based on monthly ensembles of historical sequences of observed climate (rainfall and potential evapotranspiration) that form input to a hydrological model. The outputs are probabilistic simulations of the average river flow over the forecast period (1 to 12 months ahead), at each location. The simulations are generated by the GR4J conceptual rainfall-runoff model from IRSTEA (France) calibrated on observed or naturalised flows.

3-month river flow outlook starting Apr 2020






The bar plot maps show the outlook distribution for 3, 6 and 12-month period for 64 catchments across England and Wales. Each bar plot represents the probabilistic distribution of the simulated river flow compared to the historical river flow, for the same n-month period. The probabilities fall within five categories, classified as: low, below normal, normal, above normal and high.

6-month river flow outlook starting Apr 2020



This outlook is based entirely on historical sequences and therefore does not contain any knowledge of the state of the atmosphere and ocean. It is hence possible that some of the historical sequences used might be inconsistent with current large-scale atmospheric conditions and would therefore be unlikely to occur in the next few months.



This outlook is based on monthly ensembles of historical sequences of observed climate (rainfall and potential evapotranspiration) that form input to a hydrological model. The outputs are probabilistic simulations of the average river flow over the forecast period (1 to 12 months ahead), at each location. The simulations are generated by the GR4J conceptual rainfall-runoff model from IRSTEA (France) calibrated on observed or naturalised flows.

The stack diagrams show the variation over time of the outlook distribution for a number of individual catchments. Each graph represents variation over time of the number of simulated river flows, in each month ensemble, that fall within each of seven categories: exceptionally low, notably low, below normal, normal, above normal, notably high and exceptionally high. The categories represent cumulative flow conditions, e.g. For 3-month, the simulated total 3-month flow compared to the historical 3-month flow distribution. The monthly variations can be compared to the long-term average distribution of river flows (shown as columns

on the right of each timeline graph).




This outlook is based entirely on historical sequences and therefore does not contain any knowledge of the state of the atmosphere and ocean. It is hence possible that some of the historical sequences used might be inconsistent with current large-scale atmospheric conditions and would therefore be unlikely to occur in the next few months.

Outlook based on modelled groundwater from historical climate

Period: April 2020 – March 2021

Issued on 07.04.2020 using data to the end of March

Normal conditions are expected across the UK over the next 12 months, with below normal levels possible in the latter 6 months of the year in the East of England. Exceptionally high levels are predicted in the Permo-Triassic sandstones at Skirwith for the next 5 months, remaining above normal for the remainder of the year. Note that Heathlanes has been abandoned with no levels since March 2019 and has been omitted.



This outlook is based on monthly ensembles of historical sequences of observed climate (rainfall and potential evapotranspiration) that form input to hydrological models. The outputs are probabilistic simulations of the average groundwater level over the forecast horizon (3 to 12 months ahead), at each location.

The graphs show variation over time of the number of simulated groundwater levels in each monthly ensemble,

that fall within each the seven categories: exceptionally low, notably low, below normal, normal, above normal, notably high and exceptionally high. The monthly variations can be compared to the long-term average distribution of levels, which are shown as columns on the left and right of each graph.

This outlook is based entirely on historical sequences and therefore does not contain any knowledge of the state of

the atmosphere and ocean. It is hence possible that some of the historical sequences used might be inconsistent with current large-scale atmospheric conditions and would therefore be unlikely to occur in the next few months.